

Nonrigid chiral soliton for the octet and decuplet baryons

Satoru Akiyama* and Yasuhiko Futami

Department of Physics, Tokyo University of Science, 2641, Noda, Chiba 278-8510, Japan

(Dated: February 1, 2008)

Systematic treatment of the collective rotation of the nonrigid chiral soliton is developed in the SU(3) chiral quark soliton model and applied to the octet and decuplet baryons. The strangeness degrees of freedom are treated by a simplified bound-state approach which omits the locality of the kaon wave function. Then, the flavor rotation is divided into the isospin rotation and the emission and absorption of the kaon. The kaon Hamiltonian is diagonalized by the Hartree approximation. The soliton changes the shape according to the strangeness. The baryons appear as the rotational bands of the combined system of the soliton and the kaon.

PACS numbers: 12.39.Fe, 12.38.Lg, 12.39.Ki, 14.20.Dh

I. INTRODUCTION

The soliton picture of the baryon is employed by many effective models of QCD in the low energy region: the Skyrme model [1, 2, 3, 4, 5, 6], the Nambu–Jona-Lasinio model [7, 8, 9] and the chiral quark soliton model (CQSM) [10, 11, 12, 13]. A technical feature of the picture is the separation of the internal and external degrees of freedom. The internal degrees of freedom (mesons, quarks) nonlinearly interact with each other and construct the soliton self-consistently. On the other hand, the external degrees of freedom describe the collective motions of the soliton: rotation, vibration, translational motion. To solve the coupled equations for these degrees of freedom, for technical reason, one should assume the decoupling of the internal degrees of freedom with the collective motion and employ the simplest ansatz for the soliton: the hedgehog shape. For example, in the CQSM, the equations of motion for the soliton consist of the local functions of the meson profile and the infinite eigenstates of the quarks in the background soliton. Then, the equations should be self-consistently solved. The process requires a large number of calculations by the computer. And it is hard to self-consistently incorporate the interaction between the nonrigid soliton and the collective motion. Thus, the symmetry of the soliton is a technically important ingredient to reduce the number of calculational steps. From the physical point of view, however, it seems oversimplified that one assumes the symmetry which is broken by the collective rotation.

In the previous work [14], we studied these assumption for the octet baryons with the CQSM. To overcome the complexities of the self-consistent calculation for the nonrigid soliton and incorporate the interaction between the soliton and the collective motion, we introduced the physical and/or mathematical tools: the mean field approximation for the rotated system, the tensor operator $Z_{KK_3}^{JLSI}(\hat{\mathbf{r}})$ for the grand spin which is used to expand the chiral field, and the explicit form of the isospin vector of

the chiral soliton with the quark states. The result indicated that the hedgehog shape of the soliton is stable only for the nucleon (N) and unstable for the strange baryons (Λ, Σ, Ξ). Thus, the soliton changes the shape according to the strangeness of the baryon.

Although these tools are applicable to the more general systems, the collective Hamiltonian takes the complicated form and the outlook of the collective quantization is bad due to the less symmetry of the nonrigid soliton than the hedgehog one. Therefore, in this article, we reformulate the collective Hamiltonian and improve the procedure of the collective quantization given in Ref. [14]. Furthermore, we introduce the Hartree approximation for the kaon to simplify the treatment of the particle-antiparticle creation [14]. We apply these improved tools to the octet and decuplet baryons.

In Sec. II, we review the SU(3) CQSM and the mean field approximation for the rotated system. In Sec. III, the variable transformation between the two types of the parameterization for the flavor rotation is given. In Sec. IV, first, we construct the collective Hamiltonian using the variable transformation. Next, we show the Hartree approximation for the kaon in the background soliton and define the basis for the collective quantization using the Hartree states. In Sec. V, we show the numerical results of the Hartree approximation and the collective quantization. Finally, in Sec. VI we summarize the results and discuss the relation between the deformation of the chiral soliton and the kaon.

II. MODEL AND THE MEAN FIELD APPROXIMATION FOR THE FLAVOR ROTATION

Hereafter we follow the notation used in Ref. [14] for the various quantities. The chiral quark soliton model in the case of flavor SU(3) is given by the path integral with respect to the chiral meson fields and the quark fields [10, 11, 15]. We postulate the so called cranking form [2, 16] for the meson field:

$$U^{\gamma_5}(x) = \mathcal{A}(t)B^\dagger(t)U_0^{\gamma_5}(\mathbf{r})B(t)\mathcal{A}^\dagger(t), \quad (1)$$

*Electronic address: akiyama@ph.noda.tus.ac.jp

where $U_0^{\gamma_5}(\mathbf{r})$ is the static meson field, $\mathcal{A}(t)$ describes the adiabatic rotation of the system in SU(3) flavor space, and $B(t)$ describes the spatial rotation. Then, the effective action for U^{γ_5} is reduced to

$$iS_F = N_c \log \det \left(i\partial_t + i\mathcal{A}^\dagger \dot{\mathcal{A}} + iB\dot{B}^\dagger - H' \right), \quad (2)$$

where $\det$ denotes the functional determinant for the quarks, N_c is the number of colors, and H' is the rotated quark Hamiltonian defined below.

For $U_0^{\gamma_5}$, we assume the embedding of the SU(2) field to the SU(3) matrix but do not assume the hedgehog shape [14].

We write the flavor rotation [17] as

$$\mathcal{A}(t) = \begin{pmatrix} A(t) & 0 \\ 0^\dagger & 1 \end{pmatrix} A_s(t), \quad (3)$$

where A is the flavor SU(2) rotation operator and A_s represents the rotation into the strange directions. In particular, we parameterize $A_s(t)$ as

$$A_s(t) = \exp i \begin{pmatrix} 0 & \sqrt{2}D(t) \\ \sqrt{2}D^\dagger(t) & 0 \end{pmatrix}, \quad (4)$$

where $D = (D_1, D_2)^T$ is the isodoublet spinor. From the transformation property under the flavor rotation, we call D the kaon field. In Ref. [19], it is argued that $D \sim 1/\sqrt{N_c}$ in the large N_c limit due to the Wess-Zumino term, even if the strange quark mass is light. We also employ the classification and treat D perturbatively. Although Eq. (4) was motivated by the bound-state approach [18], we will not treat the locality of the kaon wave function [19] in this article. Inclusion of the locality is a complicated task.

From the current quark mass matrix: $\hat{m} = m_0\lambda_0 + m_8\lambda_8$ ($m_3 = 0$), we obtain the rotated one:

$$\hat{m}' = \mathcal{A}^\dagger \hat{m} \mathcal{A} = m_0\lambda_0 + m_8 D_{8\mu}^{(8)}(A_s)\lambda_\mu, \quad (5)$$

where $D_{\mu\nu}^{(8)}(A_s)$ ($\mu, \nu = 1, 2, \dots, 8$) is the Wigner D function of A_s in the adjoint representation:

$$D_{\mu\nu}^{(8)}(A_s) = \frac{1}{2} \text{tr} \left(A_s^\dagger \lambda_\mu A_s \lambda_\nu \right). \quad (6)$$

The value of m_8 represents the strength of the flavor SU(3) symmetry breaking.

Here, we define the following quantities:

$$\kappa_0 \equiv 2D^\dagger D, \quad (7)$$

$$\kappa_j \equiv 2D^\dagger \tau_j D, \quad (8)$$

where $j = 1, 2, 3$, and give the explicit form of $D_{\mu\nu}^{(8)}(A_s)$ in the Appendix A. Suppose that the collective motions $\mathcal{A}$ and B are quantized and $|B\rangle$ as an eigenstate of the collective Hamiltonian. If $|B\rangle$ points to a specific direction in the isospin space,

$$\kappa_{B0} = \langle B | \kappa_0 | B \rangle, \quad (9)$$

$$\kappa_{B3} = \langle B | \kappa_3 | B \rangle \quad (10)$$

have nonzero values [14]. Then, the expectation value $\langle B | \hat{m}' | B \rangle$ may be approximated by

$$\hat{m}_B = m_0\lambda_0 + m_{B3}\lambda_3 + m_{B8}\lambda_8, \quad (11)$$

with

$$m_{B\mu} = m_8 \lim_{\kappa_{0,3} \rightarrow \kappa_{B0,3}} D_{8\mu}^{(8)}(A_s) \quad (\mu = 3, 8). \quad (12)$$

We call $\hat{m}_B$ the mean field value of $\hat{m}'$.

Using these quantities, the rotated quark Hamiltonian H' in Eq. (2) is given by

$$H' = H'_0 + \Delta H', \quad (13)$$

$$H'_0 = \frac{1}{i} \boldsymbol{\alpha} \cdot \nabla + \beta (MU_0^{\gamma_5} + \hat{m}_B), \quad (14)$$

$$\Delta H' = \beta (\hat{m}' - \hat{m}_B) = \beta T_\mu \sigma_\mu, \quad (15)$$

where M is the dynamical quark mass, σ_μ ($\mu = 1, 2, \dots, 8$) defines the fluctuation around the mean field, and $T_\mu = \lambda_\mu/2$. H'_0 contain the effects of the flavor SU(3) and SU(2) symmetry breaking through the mean field $\hat{m}_B$.

We expand S_F in power of the angular velocities ($i\mathcal{A}^\dagger \dot{\mathcal{A}}$, $iB\dot{B}^\dagger$) and $\Delta H'$ around the eigenstate of $(i\partial_t - H'_0)$ to get the effective Lagrangian $\mathcal{L}$ [12, 14]. The former corresponds to the $1/N_c$ expansion [20, 21, 22]. We retain the order of the expansion up to $O(1/N_c)$. The latter corresponds to the perturbation in power of σ_μ , which is given by the product of m_8 and the fluctuation of $D_{8\mu}^{(8)}(A_s)$ around the expectation value. We assume that the large parts of the flavor SU(3) symmetry breaking are included in the lowest order term of S_F through the mean field $\hat{m}_B$ and the residual effects can be estimated by the first order perturbation in σ_μ . In fact, the fluctuations is small, since the expectation values of $D_{8\mu}^{(8)}$ are evaluated according to the individual baryon states. For example, the fluctuation of $D_{88}^{(8)}$ starts from the difference between the $O(1/N_c)$ quantities in the large N_c limit: $-3(\kappa_0 - \kappa_{B0})/2$. Furthermore, the fluctuation is always accompanied with m_8 . The value of m_8 is about 200 MeV and fairly small compared with the typical energy scale Λ which is the energy cutoff parameter around 700 MeV. Here, the small value of m_8/Λ does not contradict the perturbative treatment of D in the large N_c limit, since the kaon field $D \sim 1/\sqrt{N_c}$ independent of the strange quark mass, as noted above.

The static meson field $U_0^{\gamma_5}$ is self-consistently determined [11] by a variation of the classical soliton energy E_{cl} which corresponds to the lowest order term of S_F . Because of Eq. (11), it introduces the isospin symmetry breaking $\kappa_{B3} \neq 0$ to the system in the body fixed frame. Here, $\kappa_{B3} \neq 0$ leads to an axially symmetric deformation of the chiral soliton [14].

It was pointed out in Ref. [27] that the Skyrmion must be improved by the flavor SU(2) rotation to resolve the "Yukawa problem." Then, the shape of the Skyrmion deviates from the hedgehog one in the body fixed frame.

The hedgehog shape of the soliton is the leading order term in the large N_c limit and the deviations correspond to the higher order terms. In our case, the effects of the rotation into the strangeness directions deform the hedgehog soliton.

III. VARIABLE TRANSFORMATION

We argue the relation between the two types of the parameterization for the flavor rotation. First, we define the local variables α and a of the rotation $\mathcal{A}$ and A , respectively by

$$\begin{aligned}\dot{\alpha}^\mu T_\mu &= -i\mathcal{A}^\dagger \dot{\mathcal{A}} \\ &= A_s^\dagger \begin{pmatrix} -iA^\dagger \dot{A} & 0 \\ 0^\dagger & 0 \end{pmatrix} A_s - iA_s^\dagger \dot{A}_s, \end{aligned} \quad (16)$$

$$\dot{a}^j \frac{\tau_j}{2} = -iA^\dagger \dot{A}, \quad (17)$$

where $\mu = 1, 2, \dots, 8$, and $j = 1, 2, 3$. Among $\dot{\alpha}^\mu$, $\dot{a}^j$, and $\dot{D}$, there is a relation:

$$\dot{\alpha}^\mu = \dot{a}^j D_{j\mu}^{(8)}(A_s) + \dot{D}^\dagger C_\mu(A_s) + C_\mu^\dagger(A_s) \dot{D}, \quad (18)$$

where C_μ is a isodoublet spinor defined by

$$C_\mu(A_s) \equiv \frac{1}{i} \text{tr} \left(A_s^\dagger \frac{\partial A_s}{\partial D^\dagger} \lambda_\mu \right), \quad (19)$$

and its explicit form is given in the Appendix A. Spinors C_μ and D are related to the Wigner D function $D_{\mu\nu}^{(8)}(A_s)$ by the following equations:

$$i \left(D^\dagger \frac{\tau_j}{2} C_\mu - C_\mu^\dagger \frac{\tau_j}{2} D \right) = \delta_{j\mu} - D_{j\mu}^{(8)}, \quad (20)$$

$$i \left(D^\dagger C_\mu - C_\mu^\dagger D \right) = \frac{2}{\sqrt{3}} \left(\delta_{8\mu} - D_{8\mu}^{(8)} \right). \quad (21)$$

Next, we investigate the relation among the canonical momenta. The effective Lagrangian $\mathcal{L}$ which is a function of $\dot{\alpha}^\mu$, is also a function of $\dot{a}^j$ and $\dot{D}$ through Eq. (18). The canonical momenta conjugate to α^μ are defined by

$$I_\mu^{\mathcal{A}} = \frac{\partial \mathcal{L}}{\partial \dot{\alpha}^\mu}. \quad (22)$$

On the other hand, the canonical momenta conjugate to a^j and D are defined by

$$I_j = \frac{\partial \mathcal{L}}{\partial \dot{a}^j} = D_{j\mu}^{(8)} I_\mu^{\mathcal{A}}, \quad (23)$$

$$P = \frac{\partial \mathcal{L}}{\partial \dot{D}^\dagger} = C_\mu I_\mu^{\mathcal{A}}. \quad (24)$$

For the isospin rotation, using Eqs. (20),(23) and (24), we obtain

$$I_j^{\mathcal{A}} = I_j + I_{Kj}, \quad (25)$$

where I_{Kj} is the isospin carried by the kaon:

$$I_{Kj} = i \left(D^\dagger \frac{\tau_j}{2} P - P^\dagger \frac{\tau_j}{2} D \right). \quad (26)$$

Furthermore, using Eq. (20), we obtain

$$Y = S - \frac{2}{\sqrt{3}} I_8^{\mathcal{A}}, \quad (27)$$

where the term $-(2/\sqrt{3})I_8^{\mathcal{A}}$ is the baryon number [26], S is the strangeness carried by kaon, and Y is the (left) hypercharge defined by

$$S = i \left(D^\dagger P - P^\dagger D \right), \quad (28)$$

$$Y = -\frac{2}{\sqrt{3}} D_{8\mu}^{(8)} I_\mu^{\mathcal{A}}. \quad (29)$$

We define

$$\tilde{P} = P - C_j I_j^{\mathcal{A}} - C_8 I_8^{\mathcal{A}}. \quad (30)$$

For the rotation into the strange direction, Eq. (24) is rewritten as

$$C_a I_a^{\mathcal{A}} = \tilde{P}. \quad (31)$$

where $a = 4, 5, 6, 7$. This equation is solved for $I_a^{\mathcal{A}}$ as follows.

$$\begin{aligned} I_4^{\mathcal{A}} &= -\frac{\sqrt{3}}{2N_1} \left(D_{84}^{(8)} F_1 + D_{85}^{(8)} G_1 \right), \\ I_5^{\mathcal{A}} &= -\frac{\sqrt{3}}{2N_1} \left(D_{85}^{(8)} F_1 - D_{84}^{(8)} G_1 \right), \\ I_6^{\mathcal{A}} &= -\frac{\sqrt{3}}{2N_2} \left(D_{86}^{(8)} F_2 + D_{87}^{(8)} G_2 \right), \\ I_7^{\mathcal{A}} &= -\frac{\sqrt{3}}{2N_2} \left(D_{87}^{(8)} F_2 - D_{86}^{(8)} G_2 \right), \end{aligned} \quad (32)$$

where N_i, F_i , and G_i ($i = 1, 2$) are functions of D and $\tilde{P}$, and given in the Appendix A.

IV. COLLECTIVE QUANTIZATION

In the same manner as Eq. (22), we introduce the canonical momenta J_j conjugate to the local variables b^j of $B(t)$ [14, 16]. Then, the collective Hamiltonian is given by

$$\begin{aligned}
\mathcal{H} &= \dot{\alpha}^\mu I_\mu^A + \dot{b}^j J_j - \mathcal{L} \\
&= E_{cl} + \frac{1}{2U_{33}} (J_3 - B_3 + \sigma_3 \Delta_{33} + \sigma_8 \Delta_{38})^2 + \sigma_3 \Gamma_3 + \sigma_8 \Gamma_8 \\
&\quad + \frac{1}{2(U_{11}V_{11} - W_{11}^2)} \sum_{j=1}^2 [V_{11} (I_j^A - \sigma_j \Delta_{11})^2 + U_{11} (J_j + \sigma_j \tilde{\Delta}_{11})^2 \\
&\quad + 2W_{11} (I_j^A - \sigma_j \Delta_{11}) (J_j + \sigma_j \tilde{\Delta}_{11})] \\
&\quad + \frac{1}{2U_{44}} [(I_4^A - \sigma_4 \Delta_{44})^2 + (I_5^A - \sigma_5 \Delta_{44})^2] + \frac{1}{2U_{66}} [(I_6^A - \sigma_6 \Delta_{66})^2 + (I_7^A - \sigma_7 \Delta_{66})^2], \tag{33}
\end{aligned}$$

where $E_{cl}, U_{\mu\nu}, V_{\mu\nu}, W_{\mu\nu}, \Delta_{\mu\nu}, \tilde{\Delta}_{\mu\nu}, B_\mu$, and Γ_μ are defined in Ref. [14] and calculated with the quark states in the background soliton. Especially, E_{cl} is the classical soliton energy. Because of the axial symmetry of the chiral field, there is the following constraint on the canonical momenta:

$$I_3^A + J_3 = 0. \tag{34}$$

In our approach, the Hamiltonian and the constraint should be considered as functions of the variables a^j, b^j, D , and the momenta I_j, J_j, P with Eqs. (25) and (32). Furthermore, we expand $\mathcal{H}$ up to the first order in power of $1/N_c$ and σ_μ . The explicit form of the collective Hamiltonian is given in Appendix B.

In Ref. [14], the derivation of the Hamiltonian from the Lagrangian was performed by the perturbation with respect to $1/N_c$ and m_8 . As a result, the Hamiltonian had a highly complicated form. On the other hand, the present form of $\mathcal{H}$ is simple and has a good foresight.

For the quantization of the system, we separate $\mathcal{H}$ as

$$\mathcal{H} = E_{cl} + \mathcal{H}_K + \mathcal{H}_{rot} + \mathcal{H}_{int}, \tag{35}$$

where $\mathcal{H}_K$ is the part containing only D and P and describes the kaon in the background soliton, $\mathcal{H}_{rot}$ is the part containing only I_j and J_j and represents the collective rotation of the system in isospin and real space, and $\mathcal{H}_{int}$ represents an interaction between the kaon and the rotation.

A. Kaon Hamiltonian $\mathcal{H}_K$

We separate $\mathcal{H}_K$ into a bilinear part $\mathcal{H}_{K0}$ and a higher order part $\mathcal{H}_{K1}$ in power of the kaon operators D and P .

$$\mathcal{H}_K = \mathcal{H}_{K0} + \mathcal{H}_{K1}. \tag{36}$$

The explicit form of the these quantities are given in Appendix B.

$\mathcal{H}_{K0}$, S [Eq. (28)], and I_{K3} [Eq. (26)] can be exactly diagonalized as follows [14, 19].

$$\mathcal{H}_{K0} = E_{ind} + \sum_{\gamma=1}^2 (\omega_\gamma \xi_\gamma^\dagger \xi_\gamma + \bar{\omega}_\gamma \bar{\xi}_\gamma^\dagger \bar{\xi}_\gamma), \tag{37}$$

$$S = \sum_{\gamma=1}^2 (\xi_\gamma^\dagger \xi_\gamma - \bar{\xi}_\gamma^\dagger \bar{\xi}_\gamma), \tag{38}$$

$$I_{K3} = \frac{1}{2} (\xi_1^\dagger \xi_1 - \bar{\xi}_1^\dagger \bar{\xi}_1 - \xi_2^\dagger \xi_2 + \bar{\xi}_2^\dagger \bar{\xi}_2), \tag{39}$$

where E_{ind} is defined in Appendix B, γ is isospin index, $\xi_\gamma^\dagger$ (ξ_γ) and $\bar{\xi}_\gamma^\dagger$ ($\bar{\xi}_\gamma$) are the creation (annihilation) operators for the kaon and antikaon, respectively, and ω_γ ($\bar{\omega}_\gamma$) is the energy eigenvalue of the kaon (antikaon). The Fock space is generated by successive operation of the creation operators on the vacuum state $|0\rangle$:

$$|n_1, \bar{n}_1, n_2, \bar{n}_2\rangle = \prod_{\gamma=1}^2 \frac{1}{\sqrt{n_\gamma! \bar{n}_\gamma!}} (\xi_\gamma^\dagger)^{n_\gamma} (\bar{\xi}_\gamma^\dagger)^{\bar{n}_\gamma} |0\rangle, \tag{40}$$

where n_γ and $\bar{n}_\gamma$ are some positive integers.

Henceforth, any quantity depending on D and P is considered as a normal ordered operator with respect to $\xi_\gamma^\dagger, \xi_\gamma, \bar{\xi}_\gamma^\dagger$, and $\bar{\xi}_\gamma$. Using this prescription, we can treat the anharmonic terms of $\mathcal{H}$ without any ambiguity. The choice of $\mathcal{H}_{K0}$ is important, because the creation and annihilation operators are given by the diagonalization of $\mathcal{H}_{K0}$. In Ref. [14], we use the $O(1)$ Hamiltonian $\mathcal{H}_0$ of $\mathcal{H}$ in the large N_c limit to define the creation and annihilation operators. The difference between $\mathcal{H}_{K0}$ and $\mathcal{H}_0$ is higher order terms in the large N_c limit. In the previous work, we classified B_3 as an $O(1/N_c)$ quantity and treated these terms in the rotation-kaon interaction. In this article, we throw this classification away and require only that $\mathcal{H}_{K0}$ contains all bilinear term with respect to D and P in $\mathcal{H}$.

To incorporate the effects of the anharmonic term $\mathcal{H}_{K1}$, we employ the Hartree approximation. In this approximation, a bare isodoublet spinor

$$X_\gamma^\dagger = \begin{pmatrix} \xi_\gamma^\dagger \\ \bar{\xi}_\gamma \end{pmatrix}, \tag{41}$$

is transformed to a dressed one $\tilde{X}_\gamma^\dagger$ by a unitary transformation:

$$\tilde{X}_\gamma^\dagger = e^{iG} X_\gamma^\dagger e^{-iG}, \quad (42)$$

where G is a Hermitian operator. Since the kaon Hamiltonian $\mathcal{H}_K$ commutes with S and I_{K3} , G takes the following form:

$$G = \sum_{\gamma=1}^2 X_\gamma^\dagger g_\gamma X_\gamma, \quad (43)$$

where g_γ are 2×2 Hermite matrices.

The unitary transformation Eq. (42) with Eq. (43) does not change S and I_{K3} . The dressed state with the good quantum numbers S and I_{K3} is defined as

$$|B(S, I_{K3})\rangle = e^{iG} |n_1, \bar{n}_1, n_2, \bar{n}_2\rangle, \quad (44)$$

where $S = n_1 - \bar{n}_1 + n_2 - \bar{n}_2$ and $I_{K3} = (n_1 - \bar{n}_1 - n_2 + \bar{n}_2)/2$. We assume that the bare state $|n_1, \bar{n}_1, n_2, \bar{n}_2\rangle$ contains only the valence (anti)kaons [19] and the kaon-antikaon pairs are created by the unitary transformation Eq. (42) with

$$g_\gamma = \begin{pmatrix} 0 & f_\gamma \\ f_\gamma^* & 0 \end{pmatrix}, \quad (45)$$

where f_γ is a complex constant. Since $S \leq 0$ for the octet and decuplet baryons,

$$|B(S, I_{K3})\rangle = e^{iG} |0, \bar{n}_1, 0, \bar{n}_2\rangle. \quad (46)$$

Here $S = -\bar{n}_1 - \bar{n}_2$ and $I_{K3} = (-\bar{n}_1 + \bar{n}_2)/2$. For the fixed S , we obtain the "multiplet" $|B(S, I_{K3})\rangle$ obeying

$$-S \geq 2I_{K3} \geq S, \quad (47)$$

since $\bar{n}_1, \bar{n}_2 \geq 0$. The energy eigenvalue of the soliton + the kaon system is calculated by

$$E_B(S, I_{K3}) = E_{cl} + \langle B(S, I_{K3}) | \mathcal{H}_K | B(S, I_{K3}) \rangle. \quad (48)$$

Since the isospin symmetry is broken by the soliton field [14], the states $|B(S, I_{K3})\rangle$ with the different I_{K3} are not degenerate even within the multiplet. Hereafter, the lowest eigenvalue is denoted by $E_{B0}(S, I_{K3})$ and the corresponding state is denoted by $|B_0(S, I_{K3})\rangle$.

The matrices g_γ are determined by the variational equations for the lowest state $|B_0(S, I_{K3})\rangle$:

$$\frac{\delta \langle B_0(S, I_{K3}) | \mathcal{H}_K | B_0(S, I_{K3}) \rangle}{\delta g_\gamma} = 0. \quad (49)$$

Thus, the values of g_γ are given for the individual $|B_0(S, I_{K3})\rangle$ with S corresponding to the baryons, and used for calculation of the higher states $|B(S, I_{K3})\rangle$ in the same multiplet.

Furthermore, we demand that the values of κ_{B0} and κ_{B3} [Eqs. (9) and (10)] are determined by the self-consistent equations for $|B_0(S, I_{K3})\rangle$:

$$\kappa_{B0} = \langle B_0(S, I_{K3}) | \kappa_0 | B_0(S, I_{K3}) \rangle, \quad (50)$$

$$\kappa_{B3} = \langle B_0(S, I_{K3}) | \kappa_3 | B_0(S, I_{K3}) \rangle. \quad (51)$$

We call the state $|B_0(S, I_{K3})\rangle$ with Eqs. (49), (50), and (51) the Hartree state.

Here, one should recognize that the Hartree state does not necessarily correspond to the absolute minimum of $E_{B0}(S, I_{K3})$ in the parameter space $(\kappa_{B0}, \kappa_{B3})$. It is because that the solution κ_{B0} of Eq. (50) is restricted by the strangeness of the baryon and the absolute minimum can be unphysical point which does not correspond to the solution.

B. Collective rotation Hamiltonian $\mathcal{H}_{rot}$ and kaon-rotation interaction Hamiltonian $\mathcal{H}_{int}$

The explicit forms of $\mathcal{H}_{rot}$ and $\mathcal{H}_{int}$ are given in Appendix B. $\mathcal{H}_{rot}$ can be diagonalized in the space spanned by the eigenstate $|J, j_3, J_3; I, i_3, I_3\rangle$, where J and J_3 (I and I_3) are the eigenvalues of the body fixed spin (isospin) operators and j_3 (i_3) is the eigenvalue of the space fixed spin (isospin) operator. The explicit representation of the eigenstate for the Euler angles is given by the direct product of the Wigner D functions in real and isospin space [16].

$\mathcal{H}_{int}$ algebraically mixes together the space of the kaon and the collective rotation. By using the constraint Eq. (34) with Eq. (25), the basis vector for the whole space is given by

$$|J, j_3, J_3; I, i_3, -(J_3 + I_{K3})\rangle |B(S, I_{K3})\rangle. \quad (52)$$

Because the total Hamiltonian $\mathcal{H}$ commutes with S , the diagonalization is well performed in the subspace with fixed eigenvalues (J, j_3) , (I, i_3) and S . However, since J_3, I_3 , and I_{K3} are not good quantum numbers individually, the rotational band of combined system of the soliton and the kaon is given by a linear combination of the basis Eq. (52):

$$|\Psi(S; J, j_3; I, i_3; \alpha)\rangle = \sum_{J_3, I_{K3}} C_{J_3, I_{K3}}^\Psi \times |J, j_3, J_3; I, i_3, -(J_3 + I_{K3})\rangle |B(S, I_{K3})\rangle, \quad (53)$$

where α indicates quantum numbers other than J, j_3, I, i_3 , and S . This means that the isospin symmetry broken by the soliton (Sec. II) would be restored by the collective rotation. One of the scales for the isospin symmetry breaking is the expectation value of κ_3 for the rotational band:

$$\kappa_{\Psi 3} = \langle \Psi | \kappa_3 | \Psi \rangle. \quad (54)$$

Of course, since we perform the quantization of the kaon and the collective rotation around a specific Hartree state $|B_0(S, I_{K3})\rangle$ which breaks the isospin symmetry, the symmetry is not perfectly restored. Thus, the calculation of the physical quantities with the rotational band Eq. (53) is an estimate.

The number of bases in Eq. (53) decreases from $(2J + 1)(2I + 1)$ due to the constraint Eq. (34) and thus the

TABLE I: The sets of parameters: the dynamical quark mass M , the current quark masses $m_u = m_d$ and m_s , and the cutoff parameter Λ . All units are MeV.

Set	M	m_u	m_s	Λ
(A)	400	6	200	700
(B)	400	6	190	700

axial symmetry of the soliton. For example, the bases for the Σ particle ($S = -1, J = 1/2, I = 1$) are given by

$$\begin{pmatrix} |J, j_3, +1/2; I, i_3, -1\rangle |B(-1, +1/2)\rangle \\ |J, j_3, -1/2; I, i_3, 0\rangle |B(-1, +1/2)\rangle \\ |J, j_3, +1/2; I, i_3, 0\rangle |B(-1, -1/2)\rangle \\ |J, j_3, -1/2; I, i_3, +1\rangle |B(-1, -1/2)\rangle \end{pmatrix}. \quad (55)$$

After the diagonalization of the total Hamiltonian $\mathcal{H}$ with these bases, we obtain the four rotational levels for Σ . We assign the level with the lowest eigenvalue as Σ in the real world. As the shape of the chiral soliton approach the hedgehog one, other higher levels become heavy and disappear from the system. For the other baryon, we equally do.

V. RESULTS

For the effective action Eq. (2), we use Schwinger's proper time regularization [8, 10, 23] with the cutoff parameter Λ . And we assume that the valence quarks are in the lowest positive energy state. Thus, we have four parameters: the dynamical quark mass M , the current quark masses $m_u = m_d$ and m_s , and Λ . We show in Table I two parameter sets that we have considered. The set (A) is the same as the parameters in Ref. [14] and used for comparison with the results. The set (B) has been obtained by fitting to the experimental values.

At first, we give the results for the set (A). Let us denote the lowest eigenvalue of $\mathcal{H}_{K0}$ [Eq. (37)] by E_{K0} . Then, $E_{cl} + E_{K0}$ is the lowest order approximation of the energy for the soliton + the kaon system. This is a sufficient quantity to see the qualitative tendency of the system in the $(\kappa_{B0}, \kappa_{B3})$ space. The more accurate calculation is performed below with the Hartree approximation. Figure 1 shows the κ_{B0} and κ_{B3} dependence of $E_{cl} + E_{K0}$ in the cases of $S = 0, -1, -2, -3$, where κ_{B0} and κ_{B3} are treated as parameters. As κ_{B0} grows, the graphs decrease for $S = -1, -2, -3$. It is because that the SU(3) symmetry breaking stabilizes the $\kappa_{B0} \neq 0$ state for the strange sectors and the $\kappa_{B0} = 0$ state for the non-strange one. On the other hand, the isospin symmetry is broken by the soliton solution. Then the $\kappa_{B3} \neq 0$ states are stable for $S = -1, -2, -3$ and the $\kappa_{B3} = 0$ state is stable for $S = 0$. However, $E_{cl} + E_{K0}$ is an even function of κ_{B3} because of the isospin symmetry of the model itself.

These graphs are flatter than our previous results [14] which represent the behavior of the lowest eigenvalue of




FIG. 1: $E_{cl} + E_{K0}$, where E_{K0} is the lowest eigenvalue of $\mathcal{H}_{K0}$ [Eq. (37)]. Here κ_{B0} and κ_{B3} are treated as parameters.

the $O(1)$ Hamiltonian of $\mathcal{H}$ [Eq. (33)] in the large N_c limit. The difference is mainly due to the terms depending on B_3 of $\mathcal{H}_{K0}$. These terms partially restore the isospin symmetry broken by the soliton in the parameter space $(\kappa_{B0}, \kappa_{B3})$. The flat regions in the graphs indicate that the Hamiltonian $E_{cl} + \mathcal{H}_{K0}$ accurately describes the combined system of kaon and soliton there.

Next, we show results of the Hartree approximation. For $S = 0, -1, -2, -3$, the Hartree states which satisfy Eqs. (49), (50), and (51) are self-consistently searched. Then, the values of κ_{B0} , κ_{B3} are determined. Table II

TABLE II: κ_{B0} , $|\kappa_{B3}|$, E_{cl} , and E_{B0} [Eq. (48)] of the Hartree states for $S = 0, -1, -2, -3$. The set of parameters is (A).

S	κ_{B0}	$ \kappa_{B3} $	E_{cl} (MeV)	E_{B0} (MeV)
0	0.00	0.00	1326	1326
-1	0.81	0.81	1463	1487
-2	1.64	1.64	1498	1600
-3	2.52	2.52	1505	1713

TABLE III: $\langle B|\kappa_0|B\rangle$, $\langle B|\kappa_3|B\rangle$, and E_B [Eq. (48)] for the multiplet $|B(S, I_{K3})\rangle$ in the Hartree approximation. $|B_0\rangle$ is the Hartree state corresponding to the lowest energy eigenvalue in a multiplet. The set of parameters is (A).

$ B(S, I_{K3})\rangle$	$\langle B \kappa_0 B\rangle$	$\langle B \kappa_3 B\rangle$	E_B (MeV)
$ B_0(0, 0)\rangle$	0.00	0.00	1326
$ B_0(-1, +1/2)\rangle$	0.81	-0.81	1487
$ B(-1, -1/2)\rangle$	0.68	0.68	1517
$ B_0(-2, +1)\rangle$	1.64	-1.64	1600
$ B(-2, 0)\rangle$	1.42	-0.22	1948
$ B(-2, -1)\rangle$	1.20	1.20	1671
$ B_0(-3, +3/2)\rangle$	2.52	-2.52	1713
$ B(-3, +1/2)\rangle$	2.27	-1.10	2301
$ B(-3, -1/2)\rangle$	2.02	0.33	2335
$ B(-3, -3/2)\rangle$	1.76	1.76	1814

shows the resultant κ_{B0} , $|\kappa_{B3}|$, E_{cl} , and E_{B0} [Eq. (48)]. It is found from the process that the stable solutions of Eq. (49) exist only in the vicinity of $\kappa_{B3} \approx 0$ or $\kappa_{B0} \approx |\kappa_{B3}|$. Moreover, Eqs. (50) and (51) restrict the choice to $\kappa_{B0} \approx |\kappa_{B3}|$. In general, $\kappa_{B0} = |\kappa_{B3}|$ is not satisfied by the dressed state Eq. (46). The only pure valence kaon state defined by Eq. (40) satisfies this relation. Our result indicates that the kaon state is stable against the particle-antiparticle creation in the vicinity of the solution. As stated in Sec. IV, the Hartree state does not correspond to the absolute minimum of $E_{B0}(S, I_{K3})$, but the energy difference is not more than 10 MeV.

The value of κ_{B3} determines the shape of the chiral soliton [14]. Since $\kappa_{B3} = 0$ for $S = 0$, the chiral soliton takes the hedgehog shape in the Hartree approximation. On the other hand, since $\kappa_{B3} \neq 0$ for $S \neq 0$, the chiral soliton takes non-hedgehog shape. Although, compared with our previous results [14], the values of κ_{B0} and $|\kappa_{B3}|$ are rather small, the conclusion about the shape of the chiral soliton does not change. From the present point of view, it is clear that the kaon states defined in Ref. [14] are unstable against the particle-antiparticle creation.

In Table III, we show the expectation values of κ_0 , κ_3 , and E_B [Eq. (48)] for each multiplet in the Hartree approximation. Here, the only multiplets which contain the Hartree state $|B_0(S, I_{K3})\rangle$ with $\kappa_{B3} \leq 0$, are listed. For $\kappa_{B3} \geq 0$, only the signs of I_{K3} and $\langle B|\kappa_0|B\rangle$ are reversed. Because of the isospin symmetry breaking, the quantities in Table III take different values even in a multiplet.

Finally, we give the results by the collective rotation. Table IV shows the possible rotational bands by the di-

TABLE IV: The possible rotational bands with Eq. (53) for the set of parameters: (A).

S	Particle	Bands (MeV)					
0	N	1382					
	Δ	1608					
-1	Λ	1205	2448				
	Σ	1239	2475	2691	5170		
	Σ^*	1453	2478	2697	4951	5173	8888
-2	Ξ	1389	1848	2109	3023		
	Ξ^*	1669	1848	2141	2764	3042	4400
-3	Ω	1738	2124	2831	3901		

TABLE V: $\kappa_{\Psi 0}$, $|\kappa_{\Psi 3}|$, Baryon masses E_Ψ . for the set of parameters: (A). Expt. denotes experimental value.

S	Particle	$\kappa_{\Psi 0}$	$ \kappa_{\Psi 3} $	E_Ψ (MeV)	Expt. (MeV)
0	N	0.00	0.00	1382	939
	Δ	0.00	0.00	1608	1232
-1	Λ	0.75	0.08	1205	1116
	Σ	0.75	0.08	1239	1193
	Σ^*	0.75	0.07	1453	1384
-2	Ξ	1.43	0.28	1389	1318
	Ξ^*	1.42	0.25	1669	1534
-3	Ω	2.13	0.30	1738	1672

agonalization with Eq. (53). As the deformations of the soliton become large, the excitation energy between the lowest and the first excited state decrease. Table V shows the lowest rotational band, which we assign as the baryon. Here, $\kappa_{\Psi 0} = \langle \Psi|\kappa_0|\Psi\rangle$ similar to Eq. (54).

The calculated masses of $S = 0$ baryons (N , Δ) are larger than the experimental value. It is pointed out in Ref. [24, 25] that these phenomena disappear by introducing the zero mode due to the hedgehog shape of the soliton. For $S = 0$, we see $\langle B|\kappa_3|B\rangle = 0$ in Table III. It means that the soliton always takes the hedgehog shape. Thus, the rotational band for $S = 0$ in Table V also produce $\kappa_{\Psi 3} = 0$ exactly.

On the other hand, for $S \neq 0$ baryons (Λ , $\Sigma^{(*)}$, $\Xi^{(*)}$, Ω), the calculated masses are light. We can recognize from a comparison between the Tables II and V, that it is due to the collective rotation in isospin and real space. The rotational bands for $S \neq 0$ contain the fluctuation caused by the transition within the multiplet $|B(S, I_{K3})\rangle$ in Table III. As a result we obtain the expectation value $\kappa_{\Psi 3} \approx 0$ in Table V.

Next, we show the possible rotational bands for the another set (B) of parameters in Tables VI.

VI. SUMMARY AND DISCUSSION

The stability of the hedgehog shape has been investigated for the octet and decuplet baryons in the chiral quark soliton model. We have expanded the collective Hamiltonian $\mathcal{H}$ up to the first order in power of $1/N_c$

TABLE VI: The possible rotational bands with Eq. (53) for the set of parameters: (B).

S	Particle	Bands (MeV)					
0	N	1382					
	Δ	1608					
-1	Λ	1178	2499				
	Σ	1212	2527	2742	5379		
	Σ^*	1425	2529	2747	5159	5380	9332
-2	Ξ	1340	1869	2122	3177		
	Ξ^*	1609	1868	2149	2925	3192	4763
-3	Ω	1654	2093	2915	4157		

and the fluctuation (σ_μ) around the mean field, and separate it into the kaon, the collective rotation, and the interaction Hamiltonian.

The kaon Hamiltonian are diagonalized by the Hartree approximation. The resultant Hartree states describe the soliton and the kaon in the background soliton. The kaons are almost in the valence states and stable against the particle-antiparticle creation. The shape of the soliton is controlled by the parameters κ_{B0} and κ_{B3} which also characterize the state of the kaon, and are self-consistently determined. The soliton takes the hedgehog shape for strangeness $S = 0$ and the non-hedgehog one for $S = -1, -2, -3$.

The Hamiltonian for the collective rotation of the soliton is diagonalized by the angular momentum basis in isospin and real space. The interaction between the rotation and the soliton is treated by the linear combination of the direct product of the angular momentum basis and the Hartree state with the constraint due to the axial symmetry of the soliton.

The $S = 0$ rotational bands include only the pure hedgehog state and have the heavy masses. Weigel *et al.* investigated the quantum correction due to the zero modes of the "hedgehog" soliton [24, 25]. The correction gives a large negative contribution (~ -400 MeV) to the N and Δ masses, and their results are in good agreement with the experimental values. Their analysis is valid also for the $S = 0$ baryons in our approach.

On the other hand, for $S \neq 0$, the rotation mixes the Hartree states with nonzero values of κ_{B3} , and the resultant rotational bands consist of the mixing of the "non-hedgehog" states. As a result, the $S \neq 0$ baryons have light masses. It is clear from the separation of the variables in the collective Hamiltonian that the light masses of the $S \neq 0$ rotational bands are due to the rotation of the non-hedgehog soliton. The calculation of Weigel *et al.* would be inapplicable to the $S \neq 0$ cases, because the soliton take the "non-hedgehog" shape and the zero mode does not arise. Thus, the consistent treatment of the deformation of the soliton and the zero mode would resolve the reversed mass order between the $S \neq 0$ and $S = 0$ baryons.

The multiplicity of the basis due to the axial symmetric deformation of the soliton necessarily produces the

excited states with the same spin-flavor quantum numbers as the ground state. For $S = -1$ baryon, the energy differences between the ground state and the first excited one are too large (~ 1 GeV) because of the small deformation of the soliton. Thus, these excited states may be beyond the validity of the model. On the other hand, the first excited states for $S = -2, -3$ have the moderate mass and would be assigned to the excited baryon in the real world. However, the strict assignment needs more symmetric treatment of the Hartree states and the collective quantization.

Our study originated from the observation [14] that the $S = -1, -2$ baryons consist of the quarks with different masses and the inertial force would deform the shape of the soliton. In fact, if SU(3) symmetry breaking vanishes ($m_s = 0$) in the quark Hamiltonian, the soliton always takes the hedgehog shape independently of the strangeness. Thus, it is surprising that Ω ($|sss\rangle$) also takes the non-hedgehog shape. The solution to the problem is in Eqs. (25) and (14). The former indicates that the emission and absorption of the kaon cause the isospin rotation of the soliton. The latter indicates that its recoil effects (the rotated quark mass) asymmetrically act on the isodoublet quarks in the body fixed frame and cause the direction of the isospin vector the asymmetry. Since the soliton consists of the valence and sea quarks, even though the masses of valence quarks are equivalent, these processes happen as far as the kaon states bear the isospin. As a result, the soliton takes the non-hedgehog shape for $S = -1, -2, -3$, stably.

Inclusion of the locality is in progress but a complicated task. Therefore we will include the results in the future work.

Acknowledgments

One of us (SA) would like to thank H. Kondo and C. Yasuda for many helpful discussions and encouragement.

APPENDIX A: FUNCTIONS RELATED TO THE ROTATION INTO THE STRANGE DIRECTION

At first, we define the following quantities:

$$\begin{aligned}
 f &= \frac{1 - \cos \sqrt{\kappa_0}}{\kappa_0}, \\
 g &= \frac{\sin \sqrt{\kappa_0}}{\sqrt{\kappa_0}}, \\
 \chi_+ &= \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \chi_- = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \\
 \tilde{\tau}_j &= \tau_j - \kappa_j f \quad (j = 1, 2, 3).
 \end{aligned}$$

The Wigner D function $D_{\mu\nu}^{(8)}(A_s)$ [Eq. (6)] are explicitly given as

where $i, j = 1, 2, 3$.

$$D_{ij}^{(8)} = \delta_{ij}(1 - \kappa_0 f) + \frac{1}{2}\kappa_i \kappa_j f^2, \quad (\text{A1})$$

$$D_{4j}^{(8)} = \frac{-ig}{\sqrt{2}} \left(D^\dagger \tilde{\tau}_j \chi_+ - \chi_+^\dagger \tilde{\tau}_j D \right), \quad (\text{A2})$$

$$D_{5j}^{(8)} = \frac{-g}{\sqrt{2}} \left(D^\dagger \tilde{\tau}_j \chi_+ + \chi_+^\dagger \tilde{\tau}_j D \right), \quad (\text{A3})$$

$$D_{6j}^{(8)} = \frac{-ig}{\sqrt{2}} \left(D^\dagger \tilde{\tau}_j \chi_- - \chi_-^\dagger \tilde{\tau}_j D \right), \quad (\text{A4})$$

$$D_{7j}^{(8)} = \frac{-g}{\sqrt{2}} \left(D^\dagger \tilde{\tau}_j \chi_- + \chi_-^\dagger \tilde{\tau}_j D \right), \quad (\text{A5})$$

$$D_{8j}^{(8)} = -\frac{\sqrt{3}}{2} g^2 \kappa_j, \quad (\text{A6})$$

$$D_{84}^{(8)} = -\sqrt{\frac{3}{2}} g(1 - \kappa_0 f) i(D_1^\dagger - D_1), \quad (\text{A7})$$

$$D_{85}^{(8)} = -\sqrt{\frac{3}{2}} g(1 - \kappa_0 f) (D_1^\dagger + D_1), \quad (\text{A8})$$

$$D_{86}^{(8)} = -\sqrt{\frac{3}{2}} g(1 - \kappa_0 f) i(D_2^\dagger - D_2), \quad (\text{A9})$$

$$D_{87}^{(8)} = -\sqrt{\frac{3}{2}} g(1 - \kappa_0 f) (D_2^\dagger + D_2), \quad (\text{A10})$$

$$D_{88}^{(8)} = 1 - \frac{3}{2} g^2 \kappa_0, \quad (\text{A11})$$

The isodoublet spinor C_μ [Eq. (19)] are represented as

$$C_j = -if(2\tau_j - \kappa_j f) D, \quad (\text{A12})$$

$$C_4 = \sqrt{2}g \left[\chi_+ + \frac{1-g}{\kappa_0 g} (D_1^\dagger + D_1) D - f(D_1^\dagger - D_1) D \right], \quad (\text{A13})$$

$$C_5 = -i\sqrt{2}g \left[\chi_+ + \frac{1-g}{\kappa_0 g} (D_1^\dagger - D_1) D - f(D_1^\dagger + D_1) D \right], \quad (\text{A14})$$

$$C_6 = \sqrt{2}g \left[\chi_- + \frac{1-g}{\kappa_0 g} (D_2^\dagger + D_2) D - f(D_2^\dagger - D_2) D \right], \quad (\text{A15})$$

$$C_7 = -i\sqrt{2}g \left[\chi_- + \frac{1-g}{\kappa_0 g} (D_2^\dagger - D_2) D - f(D_2^\dagger + D_2) D \right], \quad (\text{A16})$$

$$C_8 = -i\sqrt{3}g^2 D. \quad (\text{A17})$$

The functions N_i, F_i, G_i ($i = 1, 2$) in Eq. (32) are defined by

$$\begin{aligned} N_1 &= \left(D_{84}^{(8)} \right)^2 + \left(D_{85}^{(8)} \right)^2, & N_2 &= \left(D_{86}^{(8)} \right)^2 + \left(D_{87}^{(8)} \right)^2, \\ F_1 &= \frac{i}{2} \left[(1 + \kappa_3 f) \left(D^\dagger \tilde{P} - \tilde{P}^\dagger D \right) + (1 - \kappa_0 f) \left(D^\dagger \tau_3 \tilde{P} - \tilde{P}^\dagger \tau_3 D \right) \right], \\ F_2 &= \frac{i}{2} \left[(1 - \kappa_3 f) \left(D^\dagger \tilde{P} - \tilde{P}^\dagger D \right) - (1 - \kappa_0 f) \left(D^\dagger \tau_3 \tilde{P} - \tilde{P}^\dagger \tau_3 D \right) \right], \\ G_1 &= \frac{1}{2} (1 - \kappa_0 f) \left[\left(g - \frac{(1-g)}{\kappa_0} \kappa_3 \right) \left(D^\dagger \tilde{P} + \tilde{P}^\dagger D \right) + \left(D^\dagger \tau_3 \tilde{P} + \tilde{P}^\dagger \tau_3 D \right) \right], \\ G_2 &= \frac{1}{2} (1 - \kappa_0 f) \left[\left(g + \frac{(1-g)}{\kappa_0} \kappa_3 \right) \left(D^\dagger \tilde{P} + \tilde{P}^\dagger D \right) - \left(D^\dagger \tau_3 \tilde{P} + \tilde{P}^\dagger \tau_3 D \right) \right]. \end{aligned} \quad (\text{A18})$$

APPENDIX B: COLLECTIVE HAMILTONIAN

We define following quantities:

$$\begin{aligned}
A_{Kj} &= \frac{1}{2} (D^\dagger \tau_j P + P^\dagger \tau_j D) \quad (j = 1, 2, 3), \\
g_B &= \frac{\sin \sqrt{\kappa_{B0}}}{\sqrt{\kappa_{B0}}}, \quad \frac{1}{\Phi_{(\pm)}} = \frac{1}{2} \left(\frac{1}{U_{44}} \pm \frac{1}{U_{66}} \right), \quad \Delta_{(\pm)} = \frac{1}{2} \left(\frac{\Delta_{44}}{U_{44}} \pm \frac{\Delta_{66}}{U_{66}} \right), \\
E_{ind} &= 3m_8 g_B^2 \left[\kappa_{B0} \left(\Gamma_8 - \frac{B_3 \Delta_{38}}{U_{33}} \right) + \frac{\kappa_{B3}}{\sqrt{3}} \left(\Gamma_3 - \frac{B_3 \Delta_{33}}{U_{33}} \right) \right] + \frac{B_3^2}{2U_{33}}.
\end{aligned}$$

Furthermore, for the arbitrary three dimensional vectors $\mathbf{x} = (x_1, x_2, x_3)$ and $\mathbf{y} = (y_1, y_2, y_3)$, we introduce following symbol :

$$(\mathbf{x} \cdot \mathbf{y})_\perp = \mathbf{x} \cdot \mathbf{y} - x_3 y_3.$$

The collective Hamiltonian $\mathcal{H}$ [Eq. (33)] is expanded up to the first order in power of $1/N_c$ and σ_μ and separated as

$$\mathcal{H} = E_{cl} + \mathcal{H}_K + \mathcal{H}_{rot} + \mathcal{H}_{int}, \quad (\text{B1})$$

where E_{cl} is the classical soliton energy, $\mathcal{H}_K$ is the part containing only the kaon operators D and P and describes the kaon in the background soliton, $\mathcal{H}_{rot}$ is the part containing only the angular momenta I_j and J_j and represents the collective rotation of the system in isospin and real space, and $\mathcal{H}_{int}$ represents an interaction between the kaon and the rotation.

The kaon Hamiltonian $\mathcal{H}_K$ is given by

$$\mathcal{H}_K = \mathcal{H}_{K0} + \mathcal{H}_{K1}, \quad (\text{B2})$$

$$\begin{aligned}
\mathcal{H}_{K0} &= E_{ind} + \frac{1}{4\Phi_{(+)}} P^\dagger P + \frac{1}{4\Phi_{(-)}} P^\dagger \tau_3 P \\
&+ 3 \left[\frac{B_8^2}{8\Phi_{(+)}} + m_8 \left(\Delta_{(+)} B_8 - \Gamma_8 + \frac{B_3 \Delta_{38}}{U_{33}} \right) \right] \kappa_0 + \sqrt{3} \left(\frac{B_8}{4\Phi_{(+)}} + m_8 \Delta_{(+)} \right) S \\
&+ 3 \left[\frac{B_8^2}{8\Phi_{(-)}} + m_8 \left(\Delta_{(-)} B_8 - \frac{\Gamma_3}{\sqrt{3}} + \frac{B_3 \Delta_{33}}{\sqrt{3}U_{33}} \right) \right] \kappa_3 + \sqrt{3} \left(\frac{B_8}{4\Phi_{(-)}} + m_8 \Delta_{(-)} \right) 2I_{K3}, \quad (\text{B3})
\end{aligned}$$

$$\begin{aligned}
\mathcal{H}_{K1} &= \frac{1}{2(U_{11}V_{11} - W_{11}^2)} \left[V_{11}(\mathbf{I}_K \cdot \mathbf{I}_K)_\perp + 2\sqrt{3}m_8 (V_{11}\Delta_{11} - W_{11}\tilde{\Delta}_{11}) (\boldsymbol{\kappa} \cdot \mathbf{I}_K)_\perp \right] \\
&+ \frac{1}{\Phi_{(+)}} \left(\frac{S^2}{8} - \frac{1}{3} \mathbf{I}_K^2 \right) + \frac{I_{K3}^2}{2\Phi_{(+)}} + \frac{1}{3\Phi_{(-)}} (SI_{K3} - [\mathbf{I}_K \times \mathbf{A}_K]_3) \\
&+ \frac{\sqrt{3}}{2} \left(\frac{B_8}{4\Phi_{(-)}} + m_8 \Delta_{(-)} \right) \kappa_3 S + \sqrt{3} \left(\frac{B_8}{4\Phi_{(+)}} + m_8 \Delta_{(+)} \right) \kappa_3 I_{K3} \\
&+ \frac{\sqrt{3}}{2} \left(\frac{B_8}{4\Phi_{(+)}} - m_8 \Delta_{(+)} \right) \kappa_0 S + \left[\frac{B_8^2}{4\Phi_{(+)}} - m_8 \left(\Delta_{(+)} B_8 - \Gamma_8 + \frac{B_3 \Delta_{38}}{U_{33}} \right) \right] \kappa_0^2 \\
&+ \sqrt{3} \left(\frac{B_8}{4\Phi_{(-)}} - m_8 \Delta_{(-)} \right) \kappa_0 I_{K3} + \left[\frac{B_8^2}{4\Phi_{(-)}} - m_8 \left(\Delta_{(-)} B_8 - \frac{\Gamma_3}{\sqrt{3}} + \frac{B_3 \Delta_{33}}{\sqrt{3}U_{33}} \right) \right] \kappa_0 \kappa_3, \quad (\text{B4})
\end{aligned}$$

where $\mathcal{H}_{K0}$ is the bilinear part of $\mathcal{H}_K$ in power of D and P and $\mathcal{H}_{K1}$ is the higher order one.

The collective rotation Hamiltonian $\mathcal{H}_{rot}$ and the kaon-rotation interaction Hamiltonian $\mathcal{H}_{int}$ are given by

$$\mathcal{H}_{rot} = \frac{1}{2(U_{11}V_{11} - W_{11}^2)} [V_{11}(\mathbf{I} \cdot \mathbf{I})_{\perp} + U_{11}(\mathbf{J} \cdot \mathbf{J})_{\perp} + 2W_{11}(\mathbf{I} \cdot \mathbf{J})_{\perp}] + \frac{1}{2U_{33}}J_3^2 + \frac{J_3}{U_{33}} \left[3m_8 g_R^2 \left(\kappa_{B0}\Delta_{38} + \frac{\kappa_{B3}}{\sqrt{3}}\Delta_{33} \right) - B_3 \right], \quad (\text{B5})$$

$$\begin{aligned} \mathcal{H}_{int} = & -\frac{1}{2\Phi_{(+)}}(\mathbf{I}_K \cdot \mathbf{I})_{\perp} - \sqrt{3} \left(\frac{B_8}{4\Phi_{(+)}} + m_8\Delta_{(+)} \right) (\boldsymbol{\kappa} \cdot \mathbf{I})_{\perp} - \frac{1}{2\Phi_{(-)}} [\mathbf{I} \times \mathbf{A}_K]_3 \\ & + \frac{1}{(U_{11}V_{11} - W_{11}^2)} \left[V_{11}(\mathbf{I}_K \cdot \mathbf{I})_{\perp} + \sqrt{3}m_8 (V_{11}\Delta_{11} - W_{11}\tilde{\Delta}_{11}) (\boldsymbol{\kappa} \cdot \mathbf{I})_{\perp} \right. \\ & + W_{11}(\mathbf{I}_K \cdot \mathbf{J})_{\perp} + \sqrt{3}m_8 (W_{11}\Delta_{11} - U_{11}\tilde{\Delta}_{11}) (\boldsymbol{\kappa} \cdot \mathbf{J})_{\perp} \left. \right] \\ & + J_3 \left[\frac{S}{4\Phi_{(-)}} + \frac{I_{K3}}{2\Phi_{(+)}} + \sqrt{3} \left(\frac{B_8}{4\Phi_{(+)}} + m_8\Delta_{(+)} \right) \kappa_3 + \sqrt{3} \left(\frac{B_8}{4\Phi_{(-)}} + m_8\Delta_{(-)} \right) \kappa_0 \right. \\ & \left. - \frac{3m_8}{U_{33}} \left(\kappa_0\Delta_{38} + \frac{\kappa_3}{\sqrt{3}}\Delta_{33} \right) \right]. \quad (\text{B6}) \end{aligned}$$

-
- [1] T. H. R. Skyrme, Proc. R. Soc. London **A260**, 127 (1961).
 - [2] G. Adkin, C. Nappi, and E. Witten, Nucl. Phys. **B228**, 552 (1983).
 - [3] A. P. Balachandran, F. Lizzi, and V. G. J. Rodgers, Nucl. Phys. **B256**, 525 (1985).
 - [4] E. Guadanini, Nucl. Phys. **B236**, 35 (1984).
 - [5] P. O. Mazur, M. A. Nowak, and M. Praszalowicz, Phys. Lett. **147B**, 137,(1984).
 - [6] H. Yabu and K. Ando, Nucl. Phys. **B301**, 601 (1988).
 - [7] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961); **124**, 246 (1961).
 - [8] D. Ebert and H. Reinhardt, Nucl. Phys. **B271**, 188 (1986).
 - [9] H. Weigel, R. Alkofer, and H. Reinhardt, Nucl. Phys. **B387**, 638 (1992).
 - [10] D. I. Diakonov, V. Yu. Petrov, and P. V. Pobylitsa, Nucl. Phys. **B306**, 809 (1988).
 - [11] H. Reinhardt and R. Wünsch, Phys. Lett. B **215**, 577 (1988).
 - [12] H. Reinhardt, Nucl. Phys. **A503**, 825 (1989).
 - [13] M. Wakamatsu and H. Yoshiki, Nucl. Phys. **A524**, 561 (1991).
 - [14] S. Akiyama and Y. Futami, Phys. Rev. D **67**, 074030 (2003).
 - [15] A. Blotz, D. I. Diakonov, K. Goeke, N. W. Park, V. Yu. Petrov, and P. V. Pobylitsa, Nucl. Phys. **A555**, 765 (1993).
 - [16] E. Braaten and L. Carson, Phys. Rev. D **38**, 3525 (1988).
 - [17] D. Kaplan and I. Klebanov, Nucl. Phys. **B335**, 45 (1990).
 - [18] C. Callan and I. R. Klebanov, Nucl. Phys. **B262**, 365 (1985).
 - [19] K. M. Westerberg and I. R. Klebanov, Phys. Rev. D **50**, 5834 (1994).
 - [20] G. 't Hooft, Nucl. Phys. **B72**, 461 (1974); **B75**, 461 (1974).
 - [21] E. Witten, Nucl. Phys. **B160**, 57 (1979).
 - [22] R. Dashen, E. Jenkins, and A. V. Manohar, Phys. Rev. D **49**, 4713 (1994).
 - [23] J. Schwinger, Phys. Rev. **82**, 664 (1951).
 - [24] H. Weigel, R. Alkofer, and H. Reinhardt, Nucl. Phys. **A582**, 484 (1995).
 - [25] R. Alkofer and H. Reinhardt, *Chiral Quark Dynamics*, Springer-Verlag, Berlin, 1995.
 - [26] E. Rabinovici, A. Schwimmer, and S. Yankielowicz, Nucl. Phys. **B248**, 523 (1984).
 - [27] N. Dorey, J. Hughes, and M. P. Mattis, Phys. Rev. D **50**, 5816 (1994).